



Citation for published version:

Field, D, Bercovici, A, Berv, JS, Dunn, R, Fastovsky, D, Lyson, TR, Vajda, V & Gauthier, J 2018, 'Early Evolution of Modern Birds Structured by Global Forest Collapse at the End-Cretaceous Mass Extinction', *Current Biology*, vol. 28, no. 11, pp. 1825-1831. <https://doi.org/10.1016/j.cub.2018.04.062>

DOI:

[10.1016/j.cub.2018.04.062](https://doi.org/10.1016/j.cub.2018.04.062)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Early evolution of modern birds structured by global forest collapse at the end-Cretaceous mass extinction

Daniel J. Field^{1*}, Antoine Bercovici², Jacob S. Berv³, Regan Dunn⁴, David E. Fastovsky⁵, Tyler R. Lyson⁶, Vivi Vajda⁷, Jacques A. Gauthier⁸

¹ Milner Centre for Evolution, Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK; d.j.field@bath.ac.uk

² Department of Paleobiology MRC-121, National Museum of Natural History, Smithsonian Institution, 10th Street and Constitution Ave. NW, Washington DC 20560-0121, USA

³ Department of Ecology & Evolutionary Biology, Cornell University, 215 Tower Road, Ithaca NY, 14853, USA

⁴ Integrated Research Center, Field Museum of Natural History, 1400 S Lake Shore Dr, Chicago, IL 60605, USA

⁵ Department of Geosciences, University of Rhode Island, 9 East Alumni Ave., Kingston, RI 02881, USA

⁶ Department of Earth Sciences, Denver Museum of Nature and Science, 2001 Colorado Boulevard, Denver, CO 80205, USA

⁷ Department of Palaeobiology, Swedish Museum of Natural History, Svante Arrhenius v.9, SE-104 05 Stockholm, Sweden

⁸ Department of Geology & Geophysics, Yale University 210 Whitney Avenue, New Haven, CT, 06511, USA

*Lead Contact: d.j.field@bath.ac.uk

Summary

The fossil record and recent molecular phylogenies support an extraordinary early Cenozoic radiation of crown birds (Neornithes) following the Cretaceous-Paleogene (K-Pg) mass extinction [1-3]. However, questions remain regarding the mechanisms underlying the survival of the deepest lineages within crown birds across the K-Pg boundary, particularly since this global catastrophe eliminated even the closest stem group relatives of Neornithes [4]. Here, ancestral state reconstructions of neornithine ecology reveal a strong bias toward taxa exhibiting predominantly non-arboreal lifestyles across the K-Pg, with multiple convergent transitions toward predominantly arboreal ecologies later in the Paleocene and Eocene. By contrast, ecomorphological inferences indicate predominantly arboreal lifestyles among enantiornithines, the most diverse and widespread Mesozoic avialans [5-7]. Global paleobotanical and palynological data show that the K-Pg Chicxulub impact triggered widespread destruction of forests [8, 9]. We suggest that ecological filtering due to the temporary loss of significant plant cover across the K-Pg boundary selected against any flying dinosaurs (Avialae [10]) committed to arboreal ecologies, resulting in a predominantly non-arboreal post-extinction neornithine avifauna composed of total-clade Palaeognathae, Galloanserae, and terrestrial total-clade Neoaves that rapidly diversified into the broad range of avian ecologies familiar today. The explanation proposed here provides a unifying hypothesis for the K-Pg-associated mass extinction of arboreal stem birds, as well as for the post-K-Pg radiation of arboreal crown birds. It also provides a baseline hypothesis to be further refined pending the discovery of additional neornithine fossils from the Latest Cretaceous and earliest Paleogene.

Keywords: Mass Extinction; K-Pg; Birds; Paleobotany; Ancestral States; Ecological Selectivity

Results

Neornithine ecological selectivity across the K-Pg

Ancestral ecological reconstructions (AERs) using recent time-scaled avian phylogenies [2, 11] under likelihood, maximum parsimony, and Bayesian stochastic mapping frameworks yielded a clear ecological signal across the K-Pg boundary (Figure 1 and Supplemental Information). The deepest nodes within Neornithes are inferred to be predominantly non-arboreal, both in terms of general lifestyle and nesting substrate,

including the most recent common ancestor (MRCA) of Neornithes (the crown bird root node), and the MRCAs of Palaeognathae (ostriches and kin), Neognathae (all non-palaeognath Neornithes), and Neoaves (all neognaths excluding ducks, chickens, and their close relatives). Numerous independent transitions toward predominant arboreality are inferred for deep nodes within Neoaves early in the Cenozoic, including the extremely speciose and largely arboreal Inopinaves (a major clade of ‘core land birds’ inferred to have transitioned to predominant arboreality by 64 million years ago) [2]. These results are robust to ongoing phylogenetic uncertainty with regard to neoavian interrelationships and divergence times [1, 2, 11, 12].

Terrestrial antecedents for numerous modern arboreal bird clades

Our AERs identify several extant clades with inferred transitions to arboreality early in the Cenozoic. Some of these clades, such as Otidimorphae (turacos, cuckoos, and bustards) and Telluraves (all Inopinaves except the Hoatzin), are represented by early diverging crown clade fossils from the early Paleogene. The earliest well-constrained crown neoavian fossil (the stem mousebird *Tsidiiyazhi abini*) is inferred by Ksepka and colleagues to be predominantly arboreal [3]. However, that study also suggests that the advanced zygodactyl and semizygodactyl perching specializations of *T. abini* and other arboreal members of Telluraves are the product of multiple independent origins in the Paleocene and Eocene. Additionally, hindlimb proportions among Otidimorphae covary with degrees of arboreality and terrestriality ([13], Figure 2), and indicate that the earliest known crown otidimorph, the stem turaco *Foro panarium*, was most likely ground-dwelling, despite arboreality predominating among extant turacos.

81
82 *Assessing the extent and timeline of K–Pg forest collapse*

83 Palynological data from K–Pg boundary sections worldwide reveal a vegetation response
84 with a fern spike and floral turnover [14] , which together indicate forest destruction on a
85 global scale and a protracted (~1,000-year) onset of the recovery of climax vegetation
86 [15]. We assessed the response of forest communities to the Chicxulub impact in the
87 western interior of North America by conducting high-resolution relative abundance
88 palynological analyses (down to 1 cm sampling intervals immediately above and below
89 the boundary) of the John’s Nose K–Pg boundary section in southwestern North Dakota
90 (Figure 3). The North American K–Pg boundary sections constitute the best high-
91 resolution record of the K–Pg transition in terrestrial ecosystems [16], and John’s Nose is
92 one of two sections in southwestern North Dakota that preserve the boundary clay and
93 impact spherules, providing direct evidence of the Chicxulub impact. The palynological
94 record shows the K–Pg boundary fern spike (*Cyathidites* sp. and *Laevigatosporites* sp., >
95 70 %) 2-7 cm above the boundary, and floral turnover from 7 cm above the boundary as
96 indicated by the disappearance of typical Cretaceous pollens (K-taxa) and the dominance
97 of new pollen types in the earliest Paleocene (*Ulmipollenites krempii*, *Kurtzipites*
98 *circularis*, *K. trispissatus*, *Taxodiaceapollenites* sp., and bisaccate pollen grains; Table
99 S1).

100
101 **Discussion**

102 Synthesis of recent results from palaeontological studies and molecular divergence time
103 analyses supports a major influence of the end-Cretaceous asteroid impact on near-crown

and early crown bird evolution [1, 2, 4, 17, 18]. Fossil evidence suggests that the entirety of the neornithine stem group—including pterosaurs and all non-neornithine dinosaurs—perished in the aftermath of the impact [19], 66.02Ma [15]. This includes even the crownward-most Mesozoic avialans outside of living bird diversity, such as Ichthyornithes (*Ichthyornis* and kin), Hesperornithes (*Hesperornis* and kin), Palintropiformes (relatives of *Palintropus retusus* and *Apsaravis ukhaana*), and the diverse and globally widespread Enantiornithes (“opposite birds”), which persisted into the terminal Cretaceous (Maastrichtian) [4] (Figure 4). Although the very deepest phylogenetic divergences within Neornithes likely took place during the Mesozoic, such as the divergence between Palaeognathae and Neognathae (the neornithine root node), and that between Galloanserae and Neoaves (the deepest divergence within Neognathae), virtually the entirety of the avian crown group fossil record is restricted to sediments of Cenozoic age, and the earliest well-supported crown bird fossil is scarcely older than the end-Cretaceous, at approximately 67Ma [3, 20].

Factors influencing the post-Cretaceous survival of crown birds

Despite accumulating evidence for the dramatic influence of the Chicxulub asteroid impact on the evolutionary history of the neornithine total clade, little is known about either the factors that drove crownward stem birds such as enantiornithines to extinction, or the biological attributes of early crown birds that survived the mass extinction and radiated in its wake. Recently, selection for the toothless bill which characterizes crown birds was posited as a potential factor favoring crown bird survivorship over other contemporaneous small theropods [21]. This argument suggests that the bill would have

127 facilitated feeding on the hardy seeds and grains that may have been available as a food
128 source in the immediate aftermath of the asteroid impact. While this may be true, such a
129 scenario ignores the fact that teeth do not preclude granivory (some toothed avialans are
130 known to have fed on seeds [22]), and that a toothless bill was acquired multiple times
131 among Mesozoic Avialae, including among derived enantiornithines [23]. Additionally,
132 work incorporating fossil body size estimates, ancestral state reconstructions, and rates of
133 molecular evolution suggests that birds surviving the K–Pg mass extinction underwent
134 transient selection and filtering for reduced body size (a Lilliput Effect) [18], which may
135 have facilitated avian survival by transiently reducing their total energetic requirements.
136 Research on the evolution of neornithine breeding habits [24] suggests that, relative to
137 Enantiornithes, ancestral crown birds may have acquired proportionally larger eggs and
138 alternative nesting substrates. Moreover, evolution of a crown-grade alimentary system
139 has additionally been posited as a factor that may have influenced the post-K–Pg survival
140 of Neornithes [25]; both of these latter hypotheses are tantalizing, and warrant additional
141 research.

142 While all of the scenarios described above are plausible and not mutually
143 exclusive, none fully explain the differential survivorship of early crown birds relative to
144 crownward stem birds such as enantiornithines (which, despite differences in biological
145 attributes such as skeletal pneumaticity and growth rates, were likely biologically similar
146 to Neornithes, and probably more diverse and widespread in the terminal Cretaceous) [4].
147 Alone, these scenarios reveal an incomplete picture of broad-scale ecological selectivity
148 among the crown birds that survived the K–Pg mass extinction.

We propose a new hypothesis regarding the extinction of stem birds and the survival of crown birds across the K–Pg boundary; namely, that global deforestation caused by the Chicxulub impact induced a selective filter against the survival of arboreal birds. Given compelling evidence for transient asteroid impact-induced deforestation coincident with the end-Cretaceous mass extinction (Figure 3), the selective consequences of widespread forest destruction on i) the extinction of non-neornithine Avialae and ii) survival patterns among Neornithes must be investigated. While this hypothesis likely does not constitute the sole factor influencing end-Cretaceous avian survivorship (indeed, it is unlikely that any single hypothesis completely explains global avian survival patterns), the strength of our results, and their compatibility with other studies (e.g. [18, 21]) suggest that selection against avian arboreality across the K–Pg is likely to have played a major role in structuring the early evolutionary history of Neornithes.

Evidence for global deforestation at the K–Pg boundary

The plant fossil record and models of the effects of the Chicxulub impact provide strong evidence for the devastation of forest communities at the K–Pg boundary. Initial disruption came from energy dissipated by the impact blast, leveling trees within a radius of ~1500 km, and as intense radiated heat, which may have ignited wildfires on a global scale [26-28]. This was likely followed by acid rain resulting from the emission of sulfate-rich vapour [29], and ejection of a large quantity of soot into the atmosphere [30], potentially blocking photosynthetic activity for several years and likely inducing limited global climate cooling [31-34]. This phase of suppressed sunlight, notoriously difficult to

reconstruct, is supported by the proliferation of saprotrophs thriving on decomposing organic matter [35].

The post-impact recovery of terrestrial plant communities occurred in two phases. The first is marked by the dominance of fern spores in a ~1 cm-thick interval [36] (Figure 3). Ferns are pioneer re-colonizers of devastated landscapes, and their proliferation represents a classic example of a ‘disaster flora’ composed of taxa capable of rapidly germinating from spores and rhizomes/roots. Recent examples of fern spikes have been recorded in the recolonization of rapidly denuded landscapes, such as freshly deposited lava flows in Hawaii, slopes left barren by massive landslides induced by the 1980 eruption of Mount St. Helens, and the short-lived dominance of ferns in the Krakatau floras following its 1883 eruption [8]. The K–Pg fern spike has been identified worldwide and is an indicator of global canopy loss ([14] and references therein). Sedimentation rates based on recent high resolution radiometric dating of K–Pg bentonites from Montana [37] and the Denver Basin [15] show that establishment of the fern spike occurred within a century following the Chicxulub impact, and that the fern spike disaster flora persisted on the order of 1,000 years. This general timescale is corroborated by estimated sedimentation rates from New Zealand (Figure S1). Terrestrial floras were likely devoid of extensive closed-canopy forests during this phase.

The second phase is marked by the re-establishment of canopy vegetation: The earliest Paleocene marks a change in forest community structure compared to the Cretaceous. Typical earliest Paleocene plant assemblages are characterized by low taxonomic diversity [38–40] and by a shift of dominance toward new angiosperms and conifers (disappearance of diverse Cretaceous taxa [K-taxa], and proliferation of

Ulmipollenites krempii, *Kurtzipites* spp., palms [*Arecipites* spp.], *Taxodiaceae* and *Pinaceae*; Figure 3), long-lived plants that are indicative of modern climax communities [14, 41]. This low-diversity flora persists until the appearance of diversity hot-spots ~1.4 Myrs after the K–Pg [42].

Today, avian community diversity is negatively influenced by loss of plant diversity and habitat due to human activity including monospecific agriculture and land-use patterns [43], and the early Paleocene low-diversity floral phase may have similarly affected avian communities at that time.

Selective extinction of arboreal stem birds at the K–Pg boundary

As many as five major clades of Mesozoic non-crown avialans are known to have persisted into the final 300,000 years of the Cretaceous [4], and are inferred to have exhibited a diversity of lifestyles (Figure 4). The most diverse and globally widespread of these clades, Enantiornithes, generally exhibit feet with specializations for grasping, including long and opposable hind toes (e.g. [6, 7]). These features have long been used to identify most enantiornithines as tree-dwelling birds [5], although some enantiornithines (e.g. the Late Cretaceous taxa *Elsornis keni* and *Lectavis bretincola*) may have had reduced flight capabilities or wading specializations, suggesting a breadth of ecological habits including predominantly non-arboreal lifestyles [44]. Other crownward avialans known from the terminal Maastrichtian, such as hesperornithines and ichthyornithines [4, 45] exhibit aquatic ecologies, and the extinction of at least their marine representatives is likely related to concomitant extinctions among marine tetrapods at the K–Pg boundary [46]. Little is known regarding the ecological habits of

Palintropiformes, another clade of near-crown stem birds thought to have persisted up to the K–Pg boundary [4]. If the generally arboreal habits inferred for most Enantiornithes [5] indicate that they were largely associated with forested environments and dependent upon arboreal habitats, then the widespread destruction of forests coincident with the K–Pg transition would undoubtedly have played a major, if not absolute, role in the demise of this dominant Mesozoic clade. The same should be true for any as-yet undiscovered arboreal specialists among early crown birds and stem group ornithurines in the late Maastrichtian. More generally, a model of deforestation-related ecological selectivity across the K–Pg boundary may help explain broad-scale patterns in the early evolutionary history of other major vertebrate clades (e.g. [47]).

Selective survival of non-arboreal crown birds at the K–Pg boundary

Ancestral state reconstructions for the deepest nodes among crown birds yield a clear signal of ecological selectivity across the K–Pg boundary (Figure 1). The deepest nodes within the bird crown (the most recent common ancestor, or MRCA, of Neornithes, Palaeognathae, Neognathae, Galloanserae, and Neoaves) are unambiguously reconstructed as predominantly non-arboreal (Bayesian posterior probabilities and marginal ancestral states all = 100% across both backbone topologies). This reconstruction suggests that no lineages of arboreal crown birds crossed the K–Pg boundary, and that the numerous independent transitions toward arboreality across the neornithine tree of life—including ancient transitions within major clades such as Strisores (hummingbirds, nightjars, and kin), Otidimorphae, Columbimorphae (pigeons

and kin), and Inopinaves—took place subsequent to the K–Pg transition, presumably after global forests had rebounded from their devastation following the asteroid impact.

Evidence from the early Cenozoic neornithine fossil record

The early crown bird fossil record reveals additional support for a survivorship model whereby lineages surviving the K–Pg mass extinction were predominantly ground dwelling. For example, Otidimorphae (one of the most deeply diverging clades within Neoaves [2]) comprises a disparate group of extant birds across three major subclades: turacos (Musophagidae), cuckoos (Cuculidae), and bustards (Otididae) [2] (Figure 2). Extant turacos are medium-sized, predominantly arboreal frugivores, and bustards are large to very large obligate ground-dwellers. Cuckoos exhibit more varied ecologies, ranging from predominant ground dwelling in the ‘ground cuckoo’ clade Neomorphinae (which includes the Greater Roadrunner, *Geococcyx californianus*) to predominant arboreality in other subclades [48]. Although the early fossil record of Otidimorphae is sparse [20], the earliest known apparent crown otidimorph is the stem group turaco, *Foro panarium*, from the early Eocene of Wyoming [13]. Hindlimb proportions covary closely with ecology in Otidimorphae (Figure 2; as they do in many living birds [49]), with arboreal taxa such as turacos and arboreal cuckoos exhibiting relatively short hindlimbs, and predominantly ground-dwelling taxa such as neomorphine ground cuckoos and bustards exhibiting long hindlimbs [13]. The long hindlimb proportions of *F. panarium*, which fall within the range of small bustards, suggest that the arboreal habits of crown turacos arose from ground-dwelling ancestors, consistent with a model in which

representatives of Neoaves in the early Cenozoic retained the ground-dwelling habits of K–Pg survivors.

The earliest well-constrained neoavian fossil, *Tsidiyazhi abini*, was recently described from the early Paleocene (~62.5Ma) of New Mexico [3]. *T. abini* was inferred to represent an early stem mousebird (Coliiformes), a clade exhibiting predominantly arboreal habits today. However, ancestral state reconstructions that include *T. abini* and other early Cenozoic fossils suggest that hindlimb modifications for perching may have arisen independently in numerous arboreal clades of Telluraves, including Coliiformes, following the K–Pg mass extinction [3]. This evidence supports a model whereby early ground-dwelling neoavians repeatedly took to the trees relatively early in the Paleocene—potentially filling arboreal niches vacated by Cretaceous enantiornithines and stem ornithurines—following the recovery of global forests after the Chicxulub impact.

Conclusions

The sudden onset of the K–Pg extinction event 66.02 Ma poses significant challenges for researchers seeking to unravel its drivers (beyond the asteroid impact itself) and their consequences. These challenges are magnified for taxa with sparse K–Pg fossil records, such as birds, and for scenarios involving phenomena predicted to have taken place over ecological time intervals, such as transient yet widespread impact-related deforestation. Our picture of the contours of this extinction in terms of bird evolutionary history is still incipient; however, consilient evidence from the fossil record [3, 4, 20, 50], molecular divergence time estimates [1, 2, 11], rates of molecular evolution [18], and environmental/ecological reconstructions ([21] and this study), increasingly point to the

K–Pg impact and its consequences as playing major roles in the selective filtering of bird survivorship. All available evidence appears congruent with the globally widespread destruction of forests coincident with the K–Pg event imposing a strict filter against the persistence of avialans exhibiting arboreal ecologies.

A succession of events implicated in the destruction of global forests would have primarily affected arboreal taxa, including shock waves knocking down trees immediately following the impact, wild fires directly eliminating forest habitats, as well as reduced light levels and associated global cooling delaying forest recovery. Although a multitude of factors undoubtedly influenced avian evolution at the end-Cretaceous mass extinction, including diet [21, 25], body size [18], breeding habits [24], and flight capacity [51], selection for non-arboreal habits appears to have left an indelible mark on the early evolutionary history of crown birds, clearly discernible more than 66-million-years later.

Author Contributions:

Conceptualization, D.J.F. and J.A.G.; Methodology, D.J.F., A.B., J.S.B., R.D., T.R.L. and V.V.; Investigation, D.J.F., A.B., J.S.B. and V.V.; Writing – Original Draft, D.J.F.; Writing – Review and Editing, D.J.F., A.B., J.S.B., R.D., D.E.F., T.R.L., V.V. and J.A.G.

Acknowledgements

We thank S. Wing, A. Goswami, R. Prum, A. Heers and A. Chen for thoughtful discussion, and D.A. Pearson and M. Clark for logistical support and assistance in the field. DJF is supported by a 50th Anniversary Prize Fellowship at the University of Bath.

AB acknowledges support from the Smithsonian NMNH Deep Time Peter Buck
 Postdoctoral Fellowship. VV was supported by Swedish Research Council VR grant
 2015-4264. JSB was supported by NSF grants DGE-1650441 and DEB-1700786. JAG
 thanks the Yale Peabody Museum of Natural History. This is contribution 381 from the
 Smithsonian NMNH ETE consortium.

Declaration of Interests

The authors declare no competing interests.

References

1. Jarvis, E.D., Mirarab, S., Aberer, A.J., Li, B., Houde, P., Li, C., Ho, S.Y.W., Faircloth, B.C., Nabholz, B., Howard, J.T., et al. (2014). Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science* *346*, 1320-1331.
2. Prum, R.O., Berv, J.S., Dornburg, A., Field, D.J., Townsend, J.P., Lemmon, E.M., and Lemmon, A.R. (2015). A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature* *526*, 569-573.
3. Ksepka, D.T., Stidham, T., and Williamson, T. (2017). Early Paleocene landbird supports rapid phylogenetic and morphological diversification of crown birds after the K-Pg mass extinction. *Proceedings of the National Academy of Sciences* *114*, 8047-8052.
4. Longrich, N.R., Tokaryk, T., and Field, D.J. (2011). Mass extinction of birds at the Cretaceous-Paleogene (K-Pg) boundary. *Proceedings of the National Academy of Sciences* *108*, 15253-15257.
5. Chiappe, L.M., and Qingjin, M. (2016). *Birds of Stone: Chinese Avian Fossils from the Age of Dinosaurs*, (JHU Press).
6. Hopson, J.A. (2001). Ecomorphology of avian and nonavian theropod phalangeal proportions: Implications for the arboreal versus terrestrial origin of bird flight. *New Perspectives on the Origin and Early Evolution of Birds*, 211-235.
7. O'Connor, J.K., Chiappe, L.M., Gao, C., and Zhao, B. (2011). Anatomy of the Early Cretaceous enantiornithine bird *Rapaxavis pani*. *Acta Palaeontologica Polonica* *56*, 463-475.
8. Tschudy, R., Pillmore, C., Orth, C., Gilmore, J., and Knight, J. (1984). Disruption of the terrestrial plant ecosystem at the Cretaceous-Tertiary boundary, Western Interior. *Science* *225*, 1030-1033.

- 344 9. Vajda, V., Raine, J.I., and Hollis, C.J. (2001). Indication of global deforestation
345 at the Cretaceous-Tertiary boundary by New Zealand fern spike. *Science* 294,
346 1700-1702.
- 347 10. Gauthier, J. (1986). Saurischian monophyly and the origin of birds. *Memoirs*
348 *of the California Academy of Sciences* 8.
- 349 11. Claramunt, S., and Cracraft, J. (2015). A new time tree reveals Earth history's
350 imprint on the evolution of modern birds. *Science Advances* 1.
- 351 12. Reddy, S., Kimball, R.T., Pandey, A., Hosner, P.A., Braun, M.J., Hackett, S.J., Han,
352 K.-L., Harshman, J., Huddleston, C.J., and Kingston, S. (2017). Why do
353 phylogenomic data sets yield conflicting trees? Data type influences the avian
354 tree of life more than taxon sampling. *Systematic biology* 66, 857-879.
- 355 13. Field, D.J., and Hsiang, A.Y. (In press). A North American stem turaco, and the
356 complex biogeographic history of modern birds. *BMC Evolutionary Biology*.
- 357 14. Vajda, V., and Bercovici, A. (2014). The global vegetation pattern across the
358 Cretaceous–Paleogene mass extinction interval: A template for other
359 extinction events. *Global and Planetary Change* 122, 29-49.
- 360 15. Clyde, W.C., Ramezani, J., Johnson, K.R., Bowring, S.A., and Jones, M.M. (2016).
361 Direct high-precision U–Pb geochronology of the end-Cretaceous extinction
362 and calibration of Paleocene astronomical timescales. *Earth and Planetary*
363 *Science Letters* 452, 272-280.
- 364 16. Fastovsky, D.E., and Bercovici, A. (2016). The Hell Creek Formation and its
365 contribution to the Cretaceous–Paleogene extinction: A short primer.
366 *Cretaceous Research* 57, 368-390.
- 367 17. Feduccia, A. (1995). Explosive evolution in Tertiary birds and mammals.
368 *Science* 267, 637-638.
- 369 18. Berv, J.S., and Field, D.J. (2018). Genomic Signature of an Avian Lilliput Effect
370 across the K-Pg Extinction. *Systematic Biology* 67, 1-13.
- 371 19. Brusatte, S.L., Butler, R.J., Barrett, P.M., Carrano, M.T., Evans, D.C., Lloyd, G.T.,
372 Mannion, P.D., Norell, M.A., Peppe, D.J., and Upchurch, P. (2015). The
373 extinction of the dinosaurs. *Biological Reviews* 90, 628-642.
- 374 20. Mayr, G. (2009). *Paleogene Fossil Birds*, (Berlin: Springer).
- 375 21. Larson, D.W., Brown, C.M., and Evans, D.C. (2016). Dental disparity and
376 ecological stability in bird-like dinosaurs prior to the end-Cretaceous mass
377 extinction. *Current Biology* 26, 1325-1333.
- 378 22. Zhou, Z., and Zhang, F. (2002). A long-tailed, seed-eating bird from the Early
379 Cretaceous of China. *Nature* 418, 405-409.
- 380 23. Louchart, A., and Viriot, L. (2011). From snout to beak: the loss of teeth in
381 birds. *TRENDS in Ecology and Evolution* 26, 663-673.
- 382 24. Varricchio, D.J., and Jackson, F.D. (2016). Reproduction in Mesozoic birds and
383 evolution of the modern avian reproductive mode. *The Auk* 133, 654-684.
- 384 25. O'Connor, J., Wang, X., Sullivan, C., Wang, Y., Zheng, X., Hu, H., Zhang, X., and
385 Zhou, Z. (2018). First report of gastroliths in the Early Cretaceous basal bird
386 *Jeholornis*. *Cretaceous Research* 84, 200-208.
- 387 26. Wolbach, W.S., Gilmour, I., Anders, E., Orth, C.J., and Brooks, R.R. (1988).
388 Global fire at the Cretaceous–Tertiary boundary. *Nature* 334, 665-669.

- 389 27. Anders, E., Wolbach, W.S., and Gilmour, I. (1991). Major Wildfires at the
390 Cretaceous-Tertiary Boundary. Global biomass burning atmospheric, climatic
391 and biospheric implications, 485-492.
- 392 28. Robertson, D.S., Lewis, W.M., Sheehan, P.M., and Toon, O.B. (2013). K - Pg
393 extinction: Reevaluation of the heat - fire hypothesis. Journal of Geophysical
394 Research: Biogeosciences 118, 329-336.
- 395 29. Ohno, S., Kadono, T., Kurosawa, K., Hamura, T., Sakaiya, T., Shigemori, K.,
396 Hironaka, Y., Sano, T., Watari, T., and Otani, K. (2014). Production of sulphate-
397 rich vapour during the Chicxulub impact and implications for ocean
398 acidification. Nature Geoscience 7, 279-282.
- 399 30. Kaiho, K., Oshima, N., Adachi, K., Adachi, Y., Mizukami, T., Fujibayashi, M., and
400 Saito, R. (2016). Global climate change driven by soot at the K-Pg boundary
401 as the cause of the mass extinction. Scientific reports 6, 28427.
- 402 31. Vellekoop, J., Esmeray-Senlet, S., Miller, K.G., Browning, J.V., Sluijs, A., van de
403 Schootbrugge, B., Damsté, J.S.S., and Brinkhuis, H. (2016). Evidence for
404 Cretaceous-Paleogene boundary bolide “impact winter” conditions from New
405 Jersey, USA. Geology 44, 619-622.
- 406 32. Bardeen, C.G., Garcia, R.R., Toon, O.B., and Conley, A.J. (2017). On transient
407 climate change at the Cretaceous– Paleogene boundary due to atmospheric
408 soot injections. Proceedings of the National Academy of Sciences,
409 201708980.
- 410 33. Brugger, J., Feulner, G., and Petri, S. (2017). Baby, it's cold outside: Climate
411 model simulations of the effects of the asteroid impact at the end of the
412 Cretaceous. Geophysical Research Letters 44, 419-427.
- 413 34. Vajda, V., Ocampo, A., Ferrow, E., and Koch, C.B. (2015). Nano particles as the
414 primary cause for long-term sunlight suppression at high southern latitudes
415 following the Chicxulub impact—evidence from ejecta deposits in Belize and
416 Mexico. Gondwana Research 27, 1079-1088.
- 417 35. Vajda, V., and McLoughlin, S. (2004). Fungal proliferation at the Cretaceous-
418 Tertiary boundary. Science 303, 1489.
- 419 36. Wolfe, J.A., and Upchurch Jr, G.R. (1986). Vegetation, climatic and floral
420 changes at the Cretaceous-Tertiary boundary. Nature 324, 148.
- 421 37. Renne, P.R., Deino, A.L., Hilgen, F.J., Kuiper, K.F., Mark, D.F., Mitchell, W.S.,
422 Morgan, L.E., Mundil, R., and Smit, J. (2013). Time scales of critical events
423 around the Cretaceous-Paleogene boundary. Science 339, 684-687.
- 424 38. Johnson, K.R. (1992). Leaf-fossil evidence for extensive floral extinction at the
425 Cretaceous-Tertiary boundary, North Dakota, USA. Cretaceous Research 13,
426 91-117.
- 427 39. Johnson, K.R. (2002). Megaflora of the Hell Creek and lower Fort Union
428 Formations in the western Dakotas: Vegetational response to climate change,
429 the Cretaceous-Tertiary boundary event, and rapid marine transgression.
430 Geological Society of America Special Paper 361, 329-391.
- 431 40. Bercovici, A., Jacqueline, W., and Pearson, D. (2008). Detailed palaeontologic
432 and taphonomic techniques to reconstruct an earliest Paleocene fossil flora:

- 433 An example from southwestern North Dakota, USA. *Review of Palaeobotany*
434 *and Palynology* 151, 136-146.
- 435 41. Vajda, V., and McLoughlin, S. (2007). Extinction and recovery patterns of the
436 vegetation across the Cretaceous–Palaeogene boundary—a tool for
437 unravelling the causes of the end-Permian mass-extinction. *Review of*
438 *Palaeobotany and Palynology* 144, 99-112.
- 439 42. Johnson, K.R., and Ellis, B. (2002). A tropical rainforest in Colorado 1.4
440 million years after the Cretaceous-Tertiary boundary. *Science* 296, 2379-
441 2383.
- 442 43. Lee, M.-B., and Martin, J.A. (2017). Avian Species and Functional Diversity in
443 Agricultural Landscapes: Does Landscape Heterogeneity Matter? *PloS one* 12,
444 e0170540.
- 445 44. Chiappe, L.M., and Walker, C.A. (2002). Skeletal morphology and systematics
446 of the Cretaceous Euenantiornithes (Ornithothoraces: Enantiornithes).
447 *Mesozoic Birds: Above the Heads of Dinosaurs*, 240-267.
- 448 45. Dumont, M., Tafforeau, P., Bertin, T., Bhullar, B.-A., Field, D., Schulp, A.,
449 Strilisky, B., Thivichon-Prince, B., Viriot, L., and Louchart, A. (2016).
450 Synchrotron imaging of dentition provides insights into the biology of
451 *Hesperornis* and *Ichthyornis*, the “last” toothed birds. *BMC evolutionary*
452 *biology* 16, 178.
- 453 46. Schulte, P., Alegret, L., Arenillas, I., Arz, J.A., Barton, P.J., Bown, P.R., Bralower,
454 T.J., Christeson, G.L., Claesys, P., Cockell, C.S., et al. (2010). The Chicxulub
455 Asteroid Impact and Mass Extinction at the Cretaceous-Paleogene Boundary.
456 *Science* 327, 1214-1218.
- 457 47. Feng, Y.-J., Blackburn, D.C., Liang, D., Hillis, D.M., Wake, D.B., Cannatella, D.C.,
458 and Zhang, P. (2017). Phylogenomics reveals rapid, simultaneous
459 diversification of three major clades of Gondwanan frogs at the Cretaceous–
460 Paleogene boundary. *Proceedings of the National Academy of Sciences* 114,
461 E5864-E5870.
- 462 48. Payne, R.B. (1997). Family Cuculidae (cuckoos). In *Handbook of the Birds of*
463 *the World*, vol. 4. Sandgrouse to Cuckoos, J. del Hoyo, A. Elliott and J. Sargatal,
464 eds. (Barcelona: Lynx Edicions), pp. 508-607.
- 465 49. Norberg, U.M. (1979). Morphology of the wings, legs and tail of three
466 coniferous forest tits, the goldcrest, and the treecreeper in relation to
467 locomotor pattern and feeding station selection. *Philosophical Transactions*
468 *of the Royal Society of London B: Biological Sciences* 287, 131-165.
- 469 50. Field, D.J. (2017). Big-time insights from a tiny bird fossil. *Proceedings of the*
470 *National Academy of Sciences* 114, 7750-7752.
- 471 51. Feo, T.J., Field, D.J., and Prum, R.O. (2015). Barb geometry of asymmetrical
472 feathers reveals a transitional morphology in the evolution of avian flight.
473 *Proceedings of the Royal Society B* 282, 20142864.
- 474 52. Bercovici, A., Vajda, V., Pearson, D., Villanueva-Amadoz, U., and Kline, D.
475 (2012). Palynostratigraphy of John's Nose, a New Cretaceous—Paleogene
476 Boundary Section in Southwestern North Dakota, USA. *Palynology* 36, 36-47.

53. Field, D.J., Hanson, M., Burnham, D.A., Wilson, L.E., Super, K., Ehret, D.J., Ebersole, J.A., and Bhullar, B.A.S. (In press). Complete *Ichthyornis* skull illuminates mosaic assembly of the avian head. *Nature*.
54. Bell, A., and Chiappe, L.M. (2016). A species-level phylogeny of the Cretaceous Hesperornithiformes (Aves: Ornithuromorpha): implications for body size evolution amongst the earliest diving birds. *Journal of Systematic Palaeontology* 14, 239-251.
55. Ocampo, A., Vajda, V., and Buffetaut, E. (2006). Unravelling the Cretaceous-Paleogene (KT) catastrophe: Evidence from flora fauna and geology. *Impact Series*, 203-227.
56. Del Hoyo, J., Elliot, A., and Sargatal, J. (1992). *Handbook of the Birds of the World*. Barcelona: Lynx Editions.
57. Hsiang, A.Y., Field, D.J., Webster, T.H., Behlke, A.D.B., Davis, M.B., Racicot, R.A., and Gauthier, J.A. (2015). The origin of snakes: revealing the ecology, behavior, and evolutionary history of early snakes using genomics, phenomics, and the fossil record. *BMC Evolutionary Biology* 15, 1-22.
58. Olson, S.L. (1992). A new family of primitive landbirds from the Lower Eocene Green River Formation of Wyoming. In *Papers in avian paleontology honoring Pierce Brodkorb*, Volume 36, K.E. Campbell, ed. (Natural History Museum of Los Angeles County Science Series), pp. 137-160.
59. Team, R.D.C. (2003). *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
60. Paradis, E., Claude, J., and Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289 - 290.
61. Revell, L.J. (2012). Phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* 3, 217-223.
62. Bollback, J.P. (2006). SIMMAP: stochastic character mapping of discrete traits on phylogenies. *BMC bioinformatics* 7, 88.
63. Schliep, K.P. (2011). PHANGORN: Phylogenetic analysis using R. *Bioinformatics* 27, 592 - 593.

Figure Legends

Figure 1 Ancestral ecological reconstructions reveal bias toward non-arboreal birds across the K–Pg. Bayesian ancestral ecological reconstructions (AERs) indicate that the most deeply diverging crown bird clades, including Neornithes (all crown birds), Neognathae (Galloanserae + Neoaves), and Neoaves, were ancestrally non-arboreal ($pp > 0.99$ for each node), with numerous independent transitions toward arboreality arising in the early Cenozoic, presumably after global forests had recovered from the Chicxulub

impact. Concentric background rings demarcate geologic periods: inner grey circle at the centre is Late Cretaceous, with K–Pg boundary (66.02 Ma) shown by the red dashed line; white ring is the Paleogene (66.02–23.03Ma), separated from the Neogene (23.03–2.58Ma) by the dashed blue line. Tips extend to the present. Piecharts at nodes indicate SIMMAP posterior probabilities for ancestral ecology, under our model. Branch colors represent a single randomly sampled stochastic character map from a posterior sample of 1,000 maps. Underlying phylogeny and taxonomy follow [2]; qualitatively identical patterns are inferred using an alternative phylogenetic hypothesis [11]. See also Figures S2–S4, and doi.org/10.5281/zenodo.1204464 for alternative reconstructions.

Figure 2 Hindlimb ecomorphology suggests delayed transitions toward arboreality among Cenozoic crown birds. Early Cenozoic fossils of some modern lineages exhibiting largely arboreal habits suggest that their early antecedents were likely ground dwelling [3, 50]. Hindlimb indices ((tarsometatarsus length + tibiotarsus length)/ femur length) are plotted for the major clades within crown Otidimorphae: turacos (A; strictly arboreal), cuckoos (B; exhibiting arboreal, non-arboreal, and mixed members), and bustards (C; strictly ground-dwelling), as well as the early Eocene stem turaco *Foro panarium* (D). *F. panarium* exhibits elongated hindlimb proportions greatly exceeding the range of extant turacos and arboreal cuckoos, suggesting non-arboreal habits. Predominantly arboreal taxa are green, predominantly non-arboreal are brown, and ‘mixed’ are grey. See also Table S2.

Figure 3: Palynological record of the John's Nose section in North Dakota.

Extinction (disappearance of K-taxa) and floral turnover are evidenced by changes in relative abundance of common pollen taxa across the K–Pg boundary (modified from [52]). BC = K–Pg boundary clay. See also Figure S1 and Table S1.

Figure 4: Avialan diversity at the end-Cretaceous. At least four major clades of near-crown stem neornithines persisted into the latest Maastrichtian [4], including Enantiornithes, the most widespread and diverse clade of Mesozoic Avialae. Figure follows stratigraphic ranges from [4], with topology following recent work (e.g. [53, 54]).

STAR Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Daniel J. Field (d.j.field@bath.ac.uk).

METHOD DETAILS

Examining the extent of end-Cretaceous deforestation

Floral changes were characterized by analyzing relative abundance data of palynomorphs (pollen, spores, algae) from twelve samples spanning the K–Pg boundary from the John's Nose section, southwest North Dakota [52]. Stages of recovery were categorized based on Euclidean Distance of the relative abundance data for the 12 palynological assemblages. Raw abundance data and stratigraphic sampling resolution are presented in Table S1.

Determining the approximate timeframe of K–Pg forest recovery

The onset of fern recovery is estimated based on Iridium content in the sediments hosting the fern-spore spike based on the fall-out time calculated in Ocampo et al. [55]. The duration of fern spore dominance is based on combined calculations of sedimentation rates from K-Pg boundary successions in New Zealand [9] and the USA [15].

Neornithine ancestral ecological reconstructions

Ancestral ecological reconstructions (AERs) were performed using recent time-scaled neornithine phylogenies incorporating nearly every extant avian family-level clade [2, 11]. The 198 extant species in the original Prum *et al.* phylogenetic dataset [2], and the 229 from Claramunt and Cracraft [11] were scored as either predominantly arboreal, predominantly non-arboreal, or ‘mixed’ (for taxa that spend much of their time in both arboreal and non-arboreal settings). These alternative hypotheses span the present range of uncertainty with regard to neornithine phylogenetic topology and divergence times. Scorings were based on descriptions of general ecology, nest substrate, and foraging substrate from [56]. Discrete codings for general lifestyle and nest substrate categories are presented at doi.org/10.5281/zenodo.1204464, along with annotated R code (to reproduce all analyses) and alternative reconstructions. Categorizing behavior for the purpose of ancestral ecological reconstructions can be subjective (e.g. [57]); however, we consider our criteria for bounding categories to be consistent, and to capture the predominant ecological habits of each included species. For example, while the Greater Roadrunner (*Geococcyx californianus*) is known to nest in low trees and bushes between

1-3m above the ground [56], it is generally considered predominantly ground-dwelling (a scoring that well-reflects its general lifestyle and foraging substrate), and we therefore considered ‘non-arboreal’ to be a more suitable category for *G. californianus* than ‘mixed’. We believe that our conclusions are robust to alternative codings in similar borderline cases, such as the Shining-blue Kingfisher (*Alcedo quadibrachys*), which nests in burrows but is otherwise well classified as tree-dwelling and was therefore classified as generally arboreal. Additionally, we performed reconstructions using scores for general nest category (nests in trees vs. nests on ground vs. nests either in trees or on ground), codings that may be less subjective. These reconstructions unambiguously support ground-nesting habits as ancestral for the deepest neornithine nodes. The scenario supported by these results (clear bias toward non-arboreal nesting across the K–Pg) is qualitatively the same as that from our analysis of general lifestyle, and so are presented as supporting data (Figure S2 and doi.org/10.5281/zenodo.1204464).

Fossil bird ecomorphology

An existing morphometric dataset for Otidimorphae [13] was expanded. Digital calipers sensitive to 0.01mm were used to measure the total length of the femur, tibiotarsus, and tarsometatarsus for adult turacos, cuckoos, and bustards (raw data in Table S2). Seven species of Musophagidae were examined, representing every major musophagid subclade. Thirty species of Cuculidae were examined from across the extant diversity of cuckoos, and five species of Otidae were measured. Following [22], measurements from extant taxa were compared to measurements from the fossil stem turaco *Foro panarium* [13, 58].

QUANTIFICATION AND STATISTICAL ANALYSIS

Ancestral ecological reconstructions

Likelihood-based AERs were performed in R [59] using the `ace()` likelihood function in `ape` [60] and the `make.simmap()` Bayesian stochastic character mapping (SIMMAP) function in `phytools` [61, 62]. For both sets of analyses, we implemented a two-rate transition matrix that defines one rate for forward and reverse transitions between arboreality and non-arboreality, and another rate for forward and reverse transitions between mixed and arboreal states, and mixed and non-arboreal states (in R: `matrix(c(0, 2, 1, 2, 0, 2, 1, 2, 0), nrow=3)`). This model allows transitions that pass through the mixed phase to be favored, without preventing direct transitions. Under this model, the marginal ancestral states (from `ace`) and Bayesian posterior probabilities from SIMMAP were nearly identical ($R^2 > 0.99$), so we present our Bayesian results in Figure 1, summarized across 1,000 stochastic character maps. Predominant arboreality and predominant non-arboreality dominated inferred reconstructions throughout the tree (41.8% and 50.5% of the tree, respectively), with mixed ecologies reconstructed across only 7.7% of the tree in Figure 1. We also performed maximum parsimony reconstructions of ancestral ecological habits and nesting substrate using the `ancestral.pars()` function with regular MPR optimization implemented in the `phangorn` R package [63] (Figure S3 and doi.org/10.5281/zenodo.1204464), which corroborated the results described above.

Fossil bird ecomorphology

628 Following [13], hindlimb indices were calculated by summing the lengths of the
629 tarsometatarsus and tibiotarsus, and dividing by the length of the femur.

630

631 DATA AND SOFTWARE AVAILABILITY

632 Scripts for the ancestral ecological reconstructions, as well as the input data for these
633 analyses and all AER results are archived at doi.org/10.5281/zenodo.1204464.

634

635